

More on Solving Recurrence Relations Math 2300, Spring 2016

Second Order Linear Recurrence Relations with Constant Coefficients

Theorem: Let p_n be any particular solution to the recurrence relation $a_k = Aa_{k-1} + Ba_{k-2} + f(k)$, ignoring initial conditions. Let q_n be the general solution to the homogeneous recurrence $a_k = Aa_{k-1} + Ba_{k-2}$ again ignoring initial conditions. Then $a_n = p_n + q_n$ is the general solution to the recurrence relation $a_k = Aa_{k-1} + Ba_{k-2} + f(k)$, where the initial conditions determine the constants in q_n .

Generating Functions:

A generating function is, roughly, a polynomial that goes on forever:

$$f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots + a_nx^n + \dots$$

Definition: The generating function of a sequence $a_0, a_1, a_2, a_3, \dots$ is the expression $f(x) = a_0 + a_1x + a_2x^2 + a_3x^3 + \dots$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \dots \text{ generates } 1, 1, 1, \dots$$

$$\frac{1}{1-ax} = 1 + ax + a^2x^2 + a^3x^3 + \dots \text{ generates } 1, a, a^2, a^3, \dots$$

$$\frac{1}{(1-x)^2} = 1 + 2x + 3x^2 + 4x^3 + \dots \text{ generates } 1, 2, 3, 4, \dots$$

(The above theorem and definitions adapted from *Discrete Mathematics with Graph Theory* by Goodaire and Parmenter.)

Recursion Trees are often used for divide and conquer algorithms.

The Master Theorem gives bounds on recurrences:

The Master Theorem applies to recurrences of the following form:

$$T(n) = aT(n/b) + f(n)$$

where $a \geq 1$ and $b > 1$ are constants and $f(n)$ is an asymptotically positive function.

There are 3 cases:

1. If $f(n) = O(n^{\log_b a - \epsilon})$ for some constant $\epsilon > 0$, then $T(n) = \Theta(n^{\log_b a})$.
2. If $f(n) = \Theta(n^{\log_b a} \log^k n)$ with $k \geq 0$, then $T(n) = \Theta(n^{\log_b a} \log^{k+1} n)$.
3. If $f(n) = \Omega(n^{\log_b a + \epsilon})$ with $\epsilon > 0$, and $f(n)$ satisfies the regularity condition, then $T(n) = \Theta(f(n))$.
Regularity condition: $af(n/b) \leq cf(n)$ for some constant $c < 1$ and all sufficiently large n .

For examples of applying the Master Theorem

http://www.csanimated.com/animation.php?t=Master_theorem